

Inocor Protocol

Dilute 20ml (100mg) in 50ml of 0.9% NaCl (DO NOT mix in dextrose) to make 1.4mg/ml

Discard bag after 24 hours

Dosing (CHF): 0.75mg/kg bolus IV push over 2-3 minutes. Then start infusion of 5-10mcg/kg/min. May repeat bolus 30 minutes after start of infusion. Max daily dose is 10mg/kg (roughly 7mcg/kg/min)

Dosing chart:

Weight (kg)	0.75mg/kg (ml)	5mcg/kg/ min (ml/hr)	6mcg/kg/ min (ml/hr)	7mcg/kg/ min (ml/hr)	8mcg/kg/ min (ml/hr)	9mcg/kg/ min (ml/hr)	10mcg/kg/ min (ml/hr)
45	24.1	9.6	11.6	13.5	15.4	17.4	19.3
50	26.8	10.7	12.9	15.0	17.1	19.3	21.4
55	29.5	11.8	14.1	16.5	18.9	21.2	23.6
60	32.1	12.9	15.4	18.0	20.6	23.1	25.7
65	34.8	13.9	16.7	19.5	22.3	25.1	27.9
70	37.5	15.0	18.0	21.0	24.0	27.0	30.0
75	40.2	16.1	19.3	22.5	25.7	28.9	32.1
80	42.9	17.1	20.6	24.0	27.4	30.9	34.3
85	45.5	18.2	21.9	25.5	29.1	32.8	36.4
90	48.2	19.3	23.1	27.0	30.9	34.7	38.6
95	50.9	20.4	24.4	28.5	32.6	36.6	40.7
100	53.6	21.4	25.7	30.0	34.3	38.6	42.9
105	56.3	22.5	27.0	31.5	36.0	40.5	45.0
110	58.9	23.6	28.3	33.0	37.7	42.4	47.1
115	61.6	24.6	29.6	34.5	39.4	44.4	49.3
120	64.3	25.7	30.9	36.0	41.1	46.3	51.4
125	67.0	26.8	32.1	37.5	42.9	48.2	53.6
130	69.6	27.9	33.4	39.0	44.6	50.1	55.7

Adverse effects: Arrhythmias, nausea, vomiting, hypotension, abdominal pain, LFT elevation, anorexia, fever, chest pain, thrombocytopenia.